



**Code:** UKCAL-CWF-CON-EIA-RPT-00007-7C01

## **Volume 7C Caledonia North Appendices**

### Appendix 3-1 Marine Water Quality Baseline

Caledonia Offshore Wind Farm Ltd

5th Floor Atria One, 144 Morrison Street, Edinburgh, EH3 8EX

# Volume 7C Appendix 3-1 Marine Water Quality Baseline

|                 |                                  |
|-----------------|----------------------------------|
| <b>Code</b>     | UKCAL-CWF-CON-EIA-RPT-00007-7C01 |
| <b>Revision</b> | Issued                           |
| <b>Date</b>     | 18 October 2024                  |

# Table of Contents

|     |   |    |
|-----|---|----|
| 1   | Introduction .....                              | 1  |
| 2   | Temperature and Salinity .....                  | 3  |
| 2.1 | Overview .....                                  | 3  |
| 2.2 | Caledonia North Site .....                      | 3  |
| 2.3 | Caledonia North OECC.....                       | 6  |
| 3   | Dissolved Oxygen .....                          | 10 |
| 3.1 | Overview .....                                  | 10 |
| 3.2 | Caledonia North Site .....                      | 10 |
| 3.3 | Caledonia North OECC.....                       | 11 |
| 4   | Turbidity and Suspended Particulate Matter..... | 13 |
| 4.1 | Overview .....                                  | 13 |
| 4.2 | Caledonia North Site .....                      | 13 |
| 4.3 | Caledonia North OECC.....                       | 13 |
| 5   | pH.....   | 15 |
| 5.1 | Overview .....                                  | 15 |
| 5.2 | Caledonia North Site .....                      | 15 |
| 5.3 | Caledonia North OECC.....                       | 16 |
| 6   | References .....                                | 17 |

# List of Figures

Figure 1-1: Site-Specific Survey Locations in the Caledonia North Site and Caledonia North OECC. .... 2

Figure 2-1: Monthly mean sea surface and bottom temperature and salinity within the Caledonia North Site of the Proposed Development (Offshore) based on climatology data between 1971 and 2000. Source: Marine Directorate (2024). .... 6

Figure 2-2: Monthly mean sea surface and bottom temperature and salinity across the Caledonia North OECC based on climatology data between 1971 and 2000. Source: Marine Directorate (2024). .... 9

## List of Tables

|   |    |
|---|----|
| Table 2-1: Mean monthly sea temperature (surface and bottom) and salinity (surface and bottom) across the Caledonia North Site of the Proposed Development (Offshore) based on climatology data between 1971 and 2000. .... | 5  |
| Table 2-2: Mean monthly sea temperature (surface and bottom) and salinity (surface and bottom) across the Caledonia North OECC based on climatology data between 1971 and 2000. ....  | 8  |
| Table 3-1: Dissolved oxygen concentrations throughout the water column and across the Caledonia North Site (Source: Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)).....                               | 11 |
| Table 3-2: Dissolved oxygen concentrations throughout the water column and across the Caledonia North OECC (Source: Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).....           | 12 |
| Table 4-1: Summary of turbidity (FTU) throughout the water column and across the Caledonia North Site (Source: Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)).....                                    | 14 |
| Table 4-2: Summary of turbidity (FTU) throughout the water column and across the Caledonia North OECC (Source: Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).....                | 14 |
| Table 5-1: Summary of pH throughout the water column across the Caledonia North Site (Source: Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)). ....  | 15 |
| Table 5-2: Summary of pH throughout the water column across the Caledonia North OECC (Source: Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)). ....                                | 16 |

## Acronyms and Abbreviations

|                  |                                     |
|------------------|-------------------------------------|
| <b>CTD</b>       | Conductivity, Temperature and Depth |
| <b>MW&amp;SQ</b> | Marine Water and Sediment Quality   |
| <b>NMPI</b>      | National Marine Plan Interactive    |
| <b>OECC</b>      | Offshore Export Cable Corridor      |
| <b>OWF</b>       | Offshore Wind Farm                  |
| <b>SPM</b>       | Suspended Particulate Matter        |
| <b>SSC</b>       | Suspended Sediment Concentration    |

# 1 Introduction

1.1.1.1 This appendix covers an in-depth analysis of the baseline environment of Caledonia North in terms of marine water quality parameters in the Marine Water and Sediment Quality (MW&SQ) Study Area, such as:

- Temperature and Salinity;
- Dissolved oxygen;
- Turbidity; and
- pH.

1.1.1.2 Illustrations of the location of sample stations across the Caledonia North Site and the Caledonia North Offshore Export Cable Corridor (OECC) are shown in Figure 1-1. The results of this site-specific survey are compared with publicly available regional data, including previous Offshore Wind Farm (OWF) Environmental Impact Assessments (EIAs), to provide a robust characterisation of the entire MW&SQ Study Area.







## 2 Temperature and Salinity

### 2.1 Overview

- 2.1.1.1 Using data organised by grid cells as presented on Scotland’s National Marine Plan Interactive Map (NMPi), monthly temperature (°C) and salinity (‰) data were extracted from 20 grid cells located within the Caledonia North Site and nine grid cells within the Caledonia North OECC (Marine Directorate, 2024<sup>1</sup>). The available data layers represent a 30-year (1971-2000) temperature/salinity climatology for surface regions of the Northwest European shelf seas. From the original data, which are irregularly distributed in space and time, the mean monthly surface temperature and salinity are calculated, as well as the climatic mean annual cycle. These regional data are compared with site-specific survey data to provide a robust characterisation of temperature and salinity across the receiving environment established with high data confidence. This section refers to two measurements of salinity; parts per thousand (‰) and practical salinity units (psu). To convert salinity from parts per thousand to practical salinity units, you can use the relationship: 1 ‰ = 1 psu (e.g. 34.70 ‰ is equal to 34.70 psu).

### 2.2 Caledonia North Site

- 2.2.1.1 Salinity data was extracted from Scotland’s NMPi Map (detailed in Section 2.1) for coordinates available within the Caledonia North Site. The annual mean surface salinity (‰) was reported to remain fully marine at around 34.78‰ with minimal fluctuations throughout the year (ranging from 34.70‰ in April to 34.99‰ in October) (Table 2-1; Figure 2-1).
- 2.2.1.2 Scotland’s NMPi placed Aprils monthly average sea surface salinity within the Caledonia North Site at 34.70‰. A site-specific survey of the Caledonia North Site conducted in April observed a comparable sea surface salinity between 34.10psu and 34.60psu (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)). In general, sample stations reported stable salinity profiles across the depth of the water column, with the exception of Station ENV12, which showed a halocline (distinct vertical layer in a body of water where there is a sharp change in salinity over a relatively short depth range) between c. 10m to 25m from the surface. The development of a halocline is consistent with the freshwater input from river catchments which flow into the Moray Firth. This is validated by the fact that Station ENV12, situated in the southern part of the array, is located close to potential freshwater sources.
- 2.2.1.3 Temperature data was extracted from Scotland’s NMPi Map (detailed in Section 2.1) for coordinates available within the Caledonia North Site. Temperature values across the Caledonia North Site showed a strong seasonal signal with highs of 12.69°C in September and lows of 5.76°C in March. Moreover, during the summer months (May, June, July, August, and

September), a slight temperature variation was observed in the depth of the water column with ranges of up to 1.8°C (in June) between mean surface and bottom temperatures (Figure 2-1). This is in line with anticipated seasonal temperature stratification resulting from solar heating forming a weak thermocline (Sharples et al., 2006<sup>2</sup>; Volume 3, Chapter 2: Marine and Coastal Processes).

- 2.2.1.4 Scotland's NMPI placed April's monthly average sea surface temperature within the Caledonia North Site at 7.21°C. A site-specific survey of the Caledonia North Site conducted in April observed a comparable range of sea surface temperatures between 7.8°C and 9.7°C depending on sample location (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)). The survey output demonstrated a relatively constant sea surface temperature, suggesting a well-mixed upper layer (to approximately 10m). In addition, a clear thermocline (a distinct vertical layer within a body of water where there is a rapid change in temperature with depth) was observed with temperatures decreasing rapidly at depths ranging from 5 to 20m in all stations, with the exception of Station ENV18 and ENV34. These two outliers are located on the central seaward side of the Caledonia North Site and showed a well-mixed profile. Beyond the thermocline observed at most sample stations, temperatures were relatively stable.
- 2.2.1.5 In summary, the values reported for temperature and salinity of the Caledonia North Site are characteristic of an offshore marine environment in the North Sea throughout an annual cycle.

Table 2-1: Mean monthly sea temperature (surface and bottom) and salinity (surface and bottom) across the Caledonia North Site of the Proposed Development (Offshore) based on climatology data between 1971 and 2000.

| Month  | Sea Surface Temperature (°C) | Sea Bottom Temperature (°C) | Sea Surface Salinity (‰) | Sea Bottom Salinity (‰) |
|--|------------------------------|-----------------------------|--------------------------|-------------------------|
| January  | 7.44 (± 0.08)                | 7.71 (± 0.07)               | 34.70 (± 0.04)           | 34.76 (± 0.04)          |
| February   | 6.55 (± 0.10)                | 6.55 (± 0.07)               | 34.71 (± 0.04)           | 34.74 (± 0.03)          |
| March  | 5.76 (± 0.11)                | 5.80 (± 0.06)               | 34.72 (± 0.03)           | 34.73 (± 0.02)          |
| April  | 7.21 (± 0.02)                | 7.01 (± 0.05)               | 34.70 (± 0.05)           | 34.79 (± 0.04)          |
| May  | 8.29 (± 0.02)                | 7.55 (± 0.05)               | 34.74 (± 0.04)           | 34.85 (± 0.04)          |
| June   | 10.53 (± 0.12)               | 8.65 (± 0.15)               | 34.77 (± 0.03)           | 34.79 (± 0.03)          |
| July   | 11.35 (± 0.02)               | 10.06 (± 0.13)              | 34.83 (± 0.04)           | 34.93 (± 0.02)          |
| August   | 12.69 (± 0.10)               | 11.11 (± 0.15)              | 34.78 (± 0.02)           | 34.85 (± 0.03)          |
| September  | 12.23 (± 0.08)               | 11.85 (± 0.16)              | 34.86 (± 0.03)           | 34.93 (± 0.04)          |
| October  | 11.80 (± 0.03)               | 11.71 (± 0.06)              | 34.93 (± 0.03)           | 34.99 (± 0.04)          |
| November   | 9.93 (± 0.07)                | 9.85 (± 0.08)               | 34.82 (± 0.03)           | 34.83 (± 0.05)          |
| December   | 8.56 (± 0.07)                | 8.64 (± 0.04)               | 34.85 (± 0.02)           | 34.90 (± 0.02)          |
| Annual   | 9.36                         | 8.87                        | 34.78                    | 34.84                   |
| Note, mean (± standard deviation) are shown for each parameter (n = 20).<br>Source: Marine Directorate (2024 <sup>1</sup> ). |                              |                             |                          |                         |




Figure 2-1: Monthly mean sea surface and bottom temperature and salinity within the Caledonia North Site of the Proposed Development (Offshore) based on climatology data between 1971 and 2000. Source: Marine Directorate (2024<sup>1</sup>).

## 2.3 Caledonia North OECC

- 2.3.1.1 Salinity data was extracted from Scotland's NMPi Map (detailed in Section 2.1) for all coordinates available within the Caledonia North OECC. The annual mean surface salinity (‰) was reported to remain fully marine at around 34.74‰ with minimal fluctuations throughout the year (ranging from 34.63‰ in April to 34.89 ‰ in October) (Table 2-2). A slight variation in salinity was recorded in the depth of the water column where the annual average reported a variance of 0.06‰ favouring a slightly more saline environment at the sea bottom.
- 2.3.1.2 Scotland's NMPi placed April's monthly average sea surface salinity within Caledonia North OECC at 34.70‰. A site-specific survey of the Caledonia North OECC conducted in April observed a comparable sea surface salinity between 33.8psu and 34.40psu (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)). Sampled stations within the Caledonia North OECC reported slightly decreased salinity in surface waters. This reduced salinity in surface waters, and therefore development of a halocline, is consistent with the freshwater input from river catchments which flow into the Moray Firth.
- 2.3.1.3 Temperature data was extracted from Scotland's NMPi Map for all coordinates available within the Caledonia North OECC. Temperature values exhibited a strong seasonal signal with highs of 12.88°C in September and lows of 5.58°C in March. Moreover, during the summer months (May, June, July, and

August), a slight temperature variation was observed in the depth of the water column with variations of up to 2.04°C in June between mean surface and mean bottom temperatures (Figure 2-2:). This is in line with anticipated seasonal temperature stratification resulting from solar heating forming a weak thermocline.

- 2.3.1.4 Scotland's NMPI placed April's monthly average sea surface temperature within the Caledonia North OECC at 7.17°C. A site-specific survey of the Caledonia North OECC conducted in April observed a comparable range of sea surface temperatures between 6.9°C and 8.2°C depending on the measurement location (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)). The survey data reported a general increase in sea surface temperature associated with distance from shore, where the nearshore Station ENV17 recorded a temperature of 6.9°C which increased to 8.1°C at the furthest offshore Station ENV36. In addition, a clear thermocline was observed at depths ranging from approximately 10 to 15m in the offshore environment, whereas nearshore stations recorded a well-mixed profile throughout. Reports observed a narrowing temperature range (approximately 0.5°C) beyond depths of 30m which continued until temperature remained relatively stable at 40m and beyond (ranging from 7.2°C to 7.6°C).
- 2.3.1.5 In summary, these values reported for temperature and salinity of the Caledonia North OECC are characteristic of a typical coastal environment throughout an annual cycle with inputs from freshwater sources.

Table 2-2: Mean monthly sea temperature (surface and bottom) and salinity (surface and bottom) across the Caledonia North OECC based on climatology data between 1971 and 2000.

| Month   | Sea Surface Temperature (°C) | Sea Bottom Temperature (°C) | Sea Surface Salinity (‰) | Sea Bottom Salinity (‰) |
|---|------------------------------|-----------------------------|--------------------------|-------------------------|
| January   | 7.29 (± 0.03)                | 7.59 (± 0.04)               | 34.64 (± 0.04)           | 34.71 (± 0.02)          |
| February  | 6.39 (± 0.05)                | 6.42 (± 0.04)               | 34.66 (± 0.03)           | 34.69 (± 0.02)          |
| March   | 5.58 (± 0.05)                | 5.71 (± 0.01)               | 34.68 (± 0.02)           | 34.69 (± 0.01)          |
| April   | 7.17 (± 0.02)                | 6.92 (± 0.03)               | 34.63 (± 0.03)           | 34.74 (± 0.02)          |
| May   | 8.32 (± 0.05)                | 7.47 (± 0.02)               | 34.69 (± 0.02)           | 34.78 (± 0.02)          |
| June  | 10.73 (± 0.14)               | 8.69 (± 0.15)               | 34.72 (± 0.02)           | 34.75 (± 0.02)          |
| July  | 11.40 (± 0.03)               | 10.13 (± 0.10)              | 34.78 (± 0.05)           | 34.90 (± 0.01)          |
| August  | 12.88 (± 0.05)               | 11.24 (± 0.11)              | 34.75 (± 0.01)           | 34.81 (± 0.02)          |
| September   | 12.33 (± 0.07)               | 12.01 (± 0.11)              | 34.82 (± 0.02)           | 34.89 (± 0.03)          |
| October   | 11.82 (± 0.00)               | 11.75 (± 0.02)              | 34.89 (± 0.03)           | 34.95 (± 0.03)          |
| November  | 9.79 (± 0.04)                | 9.71 (± 0.05)               | 34.76 (± 0.03)           | 34.75 (± 0.04)          |
| December  | 8.49 (± 0.07)                | 8.65 (± 0.04)               | 34.83 (± 0.02)           | 34.88 (± 0.02)          |
| Annual Average  | 9.35 (± 2.37)                | 8.86 (± 2.03)               | 34.74 (± 0.08)           | 34.80 (± 0.08)          |
| Note, mean (± standard deviation) are shown for each parameter (n = 9).<br>Source: Marine Directorate (2024 <sup>1</sup> ). |                              |                             |                          |                         |



Figure 2-2: Monthly mean sea surface and bottom temperature and salinity across the Caledonia North OECC based on climatology data between 1971 and 2000. Source: Marine Directorate (2024<sup>1</sup>).



## 3 Dissolved Oxygen

### 3.1 Overview

- 3.1.1.1 Dissolved oxygen levels are indicative of oxygen sufficiency, with concentrations below 6mg/l considered oxygen deficient and below 2mg/l as hypoxic. Typical dissolved oxygen concentrations in the North Sea range from 6mg/l to 10mg/l (Mahaffey *et al.*, 2020<sup>3</sup>). In the OSPAR Quality Status Report (2023<sup>4</sup>), a DO concentration of >6mg/l near the seafloor is used as an indicator of a healthy marine environment.

### 3.2 Caledonia North Site

- 3.2.1.1 Dissolved oxygen was recorded throughout the water column for four locations across the Caledonia North Site associated with CTD profiling. As a result, measured dissolved oxygen concentrations are available as profiles through the water column, with a summary for each sample station presented in Table 3-1. In the OSPAR Intermediate Assessment (OSPAR, 2017<sup>5</sup>), a dissolved oxygen concentration of >6mg/l near the seafloor is used as an indicator of a healthy marine environment. All dissolved oxygen concentrations were >6mg/l throughout the water column for each sampled station within the Caledonia North Site, based on the minimum and maximum levels and water column profiles (Table 3-1).
- 3.2.1.2 Results indicate that concentrations of dissolved oxygen ranged from a minimum of 9.4 mg/l to a maximum of 10.4 mg/l. The highest dissolved oxygen concentrations were reported in surface samples within the top 10 m of the water column. This is consistent with literature that suggests surface mixing contributes to heightened air-sea gas exchange. These data are characteristic of a typical offshore coastal region of the North Sea (Mahaffey *et al.*, 2020<sup>3</sup>). The observed decline of dissolved oxygen throughout the water column may be due to a variety of physical (e.g., stratification and mixing) and/or biological factors (e.g., variations in production via photosynthesis and respiratory consumption by a wide range of organisms).

Table 3-1: Dissolved oxygen concentrations throughout the water column and across the Caledonia North Site (Source: Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area))

| Sample Station  | Range of Dissolved Oxygen (mg/l) |      |         | Dissolved oxygen (mg/l) at Specific Depth |      |           | Station Depth (m LAT)* |
|---|----------------------------------|------|---------|---|------|-----------|------------------------|
|   | Min                              | Max  | Surface | 25m                                       | 50m  | Max Depth |                        |
| ENV07   | 9.4                              | 10.0 | 10.0    | 9.5                                       | 9.4  | 9.4       | 53                     |
| ENV18   | 9.9                              | 10.1 | 10.1    | 9.9                                       | 9.9  | 9.9       | 54                     |
| ENV33   | 10.0                             | 10.3 | 10.3    | 10.1                                      | 10.0 | 10.0      | 54                     |
| ENV35   | 9.8                              | 10.4 | 10.4    | 9.9                                       | 9.9  | 9.9       | 54                     |
| * Note, the maximum station depth is consistent across all sample stations for dissolved oxygen measurements taken in the Caledonia North Site. |                                  |      |         |   |      |           |                        |

### 3.3 Caledonia North OECC

- 3.3.1.1 Dissolved oxygen was recorded throughout the water column for eight locations across the Caledonia North OECC associated with the CTD profiling. As a result, measured dissolved oxygen concentrations are available as profiles through the water column, with a summary for each sample station presented in Table 3-2. All dissolved oxygen concentrations were >6mg/l throughout the water column for each samples station within the Caledonia North OECC (Table 3-2), indicating a healthy marine environment (OSPAR, 2017<sup>5</sup>).
- 3.3.1.2 Survey results showed concentrations of dissolved oxygen range from a minimum of 9.0 mg/l to a maximum of 10.7 mg/l. The highest dissolved oxygen concentrations were reported in surface samples within the top 5 m to 10 m of the water column. This is consistent with literature that suggests surface mixing contributes to air-sea gas exchange and corresponds to the fresher inputs recorded in the salinity profiles. These data are characteristic of typical dissolved oxygen concentrations in offshore coastal region of the North Sea (Mahaffey *et al.*, 2020<sup>3</sup>). The observed decline of dissolved oxygen throughout the water column may be due to a variety of physical (such as gas exchange with the atmosphere, stratification and mixing) and/ or biological factors (such as variations in production via photosynthesis and respiratory consumption by a wide range of organisms).

Table 3-2: Dissolved oxygen concentrations throughout the water column and across the Caledonia North OECC (Source: Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).

| Sample Station  | Range of Dissolved Oxygen (mg/l) |      |         | Dissolved Oxygen at Specific Depth (mg/l) |     |           | Station Depth (m LAT)* |
|---|----------------------------------|------|---------|---|-----|-----------|------------------------|
|   | Min                              | Max  | Surface | 25m                                       | 50m | Max Depth |                        |
| ENV53   | 9.4                              | 9.6  | 9.6     | N/A                                       | N/A | 9.5       | 28                     |
| ENV17   | 9.0                              | 10.4 | 10.3    | 9.4                                       | 9.1 | 9.1       | 73                     |
| ENV26   | 9.0                              | 10.2 | 10.2    | 9.5                                       | 9.1 | 9.0       | 109                    |
| ENV04   | 9.1                              | 10.6 | 10.6    | 9.6                                       | 9.2 | 9.1       | 81                     |
| ENV13   | 9.0                              | 10.4 | 10.3    | 9.5                                       | 9.1 | 9.0       | 62                     |
| ENV29   | 9.2                              | 10.4 | 10.4    | 9.6                                       | 9.3 | 9.2       | 81                     |
| ENV31   | 9.0                              | 10.2 | 10.2    | 9.5                                       | 9.2 | 9.1       | 73                     |
| ENV36   | 9.0                              | 10.7 | 10.7    | 9.5                                       | 9.3 | 9.0       | 71                     |
| * Note, the maximum station depth is consistent across all sample stations for dissolved oxygen measurements taken in the Caledonia North OECC. |                                  |      |         |   |     |           |                        |

## 4 Turbidity and Suspended Particulate Matter

### 4.1 Overview

- 4.1.1.1 Suspended Particulate Matter (SPM) concentrations differ between the northern and southern regions of the North Sea. The northern North Sea with weaker tidal currents and deeper waters, has lower SPM concentrations compared to the southern region. Baseline SPM values in the Study Area are generally low (<5mg/l) (United Kingdom Marine Monitoring and Assessment Strategy, 2018<sup>6</sup>). However, storm events can temporarily increase near seabed SSC due to wave action. Coarser sediments are transported short distances before redeposition, while finer materials are carried further in the direction of net tidal flow (Cefas, 2016<sup>7</sup>).
- 4.1.1.2 Turbidity in the North Sea varies by location, season, weather and human activities. Typical turbidity levels in the North Sea are generally low to moderate, where coastal or estuarine environments are more turbid. In the relatively clear offshore waters of the North Sea, turbidity levels may range from around 1 to 5 Formazan Turbidity Units (FTU) under normal conditions. However, turbidity levels can be higher in nearshore areas, especially in areas influenced by river runoff, coastal erosion, or industrial activities.

### 4.2 Caledonia North Site

- 4.2.1.1 A site-specific survey (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)) measured the turbidity of the Caledonia North Site in FTU, results of which are presented in Table 4-1. Overall, turbidity profiles revealed that the water turbidity remained relatively consistent throughout the water column.

### 4.3 Caledonia North OECC

- 4.3.1.1 A site-specific survey (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)) measured the turbidity within the Caledonia North OECC in FTU, the results of which are presented in Table 4-2. Overall, turbidity profiles indicate that the water turbidity was relatively consistent throughout the water column. A slight increase in turbidity, together with data spikes, were observed in the deepest 5m to 10m at most stations. The data spikes are consistent with the instrument sensor becoming momentarily obscured by suspended particles, debris or fauna in the water column. For example, the nearshore Station ENV53 generally exhibited more turbidity than the offshore stations, which may be attributed to the spring plankton bloom observed in the seabed imagery at this station (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).

Table 4-1: Summary of turbidity (FTU) throughout the water column and across the Caledonia North Site (Source: Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)).

| Sample Station | Range of Turbidity (FTU) |     |         | Turbidity (FTU) at Specific Depth |     |           |
|----------------|--------------------------|-----|---------|-----------------------------------|-----|-----------|
|                | Min                      | Max | Surface | 25m                               | 50m | Max Depth |
| ENV07          | 1.2                      | 5.1 | 1.4     | 1.3                               | 1.4 | 1.4       |
| ENV18          | 1.3                      | 5.4 | 1.5     | 1.5                               | 1.6 | 1.8       |
| ENV33          | 1.3                      | 8.3 | 1.5     | 1.7                               | 1.8 | 1.6       |
| ENV35          | 1.4                      | 5.2 | 1.5     | 1.6                               | 1.8 | 1.6       |

Table 4-2: Summary of turbidity (FTU) throughout the water column and across the Caledonia North OECC (Source: Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).

| Sample Station | Range of Turbidity (FTU) |     |         | Turbidity (FTU) at Specific Depth |     |           |
|----------------|--------------------------|-----|---------|-----------------------------------|-----|-----------|
|                | Min                      | Max | Surface | 25m                               | 50m | Max Depth |
| ENV53          | 1.8                      | 6.7 | 2.1     | N/A                               | N/A | 2.7       |
| ENV17          | 1.3                      | 4.1 | 1.6     | 1.5                               | 1.8 | 2.0       |
| ENV26          | 1.4                      | 2.8 | 1.5     | 1.5                               | 1.6 | 2.2       |
| ENV04          | 1.3                      | 3.3 | 1.5     | 1.5                               | 1.8 | 2.4       |
| ENV13          | 1.4                      | 4.3 | 1.5     | 1.6                               | 2.2 | 2.2       |
| ENV29          | 1.4                      | 2.7 | 1.5     | 1.5                               | 1.5 | 2.1       |
| ENV31          | 1.4                      | 3.4 | 1.6     | 1.6                               | 1.8 | 2.4       |
| ENV36          | 1.2                      | 2.7 | 1.7     | 1.4                               | 1.3 | 2.5       |

## 5 pH

### 5.1 Overview

- 5.1.1.1 Fluctuations in pH can indicate changes in water chemistry, such as inputs of acid or alkaline substances from natural sources or human activities (such as coastal runoff). Monitoring pH helps assess the health and stability of marine ecosystems and identify potential sources of pollution or environmental stress. Overall, maintaining stable pH levels in seawater is essential for supporting healthy marine ecosystems, regulating biogeochemical cycles, and sustaining valuable ecosystem services provided by the oceans. The normal pH of seawater in the Moray Firth can vary depending on factors such as location, season, weather conditions, and local environmental influences.
- 5.1.1.2 Data extracted from Copernicus for pH levels between 2021 and 2024 (Copernicus, 2024<sup>8</sup>), focusing on a central data point within the Caledonia North Site, indicates an annual average pH of 8.06. The typical pH of seawater in the Moray Firth varies based on location, season, and environmental influences. Air-sea interaction with carbon dioxide generally keeps seawater pH within a range of 7.5 to 8.5, aligning with the survey results.

### 5.2 Caledonia North Site

- 5.2.1.1 A site-specific survey measured pH levels at various depths in the water column across the Caledonia North Site (Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)), results of which are displayed in Table 5-1. The pH values generally remained stable throughout the water column and across the Caledonia North Site, with a pH range of 8.3 to 8.

Table 5-1: Summary of pH throughout the water column across the Caledonia North Site (Source: Volume 7B, Appendix 4-1: Environmental Baseline Report (Array Area)).

| Sample Station | Range of pH |     |         | pH at Specific Depth |     |           |
|----------------|-------------|-----|---------|----------------------|-----|-----------|
|                | Min         | Max | Surface | 25m                  | 50m | Max Depth |
| ENV07          | 8.0         | 8.0 | 8.0     | 8.0                  | 8.0 | 8.0       |
| ENV18          | 8.2         | 8.3 | 8.2     | 8.2                  | 8.3 | 8.3       |
| ENV33          | 8.0         | 8.0 | 8.0     | 8.0                  | 8.0 | 8.0       |
| ENV35          | 8.0         | 8.0 | 8.0     | 8.0                  | 8.0 | 8.0       |

## 5.3 Caledonia North OECC

5.3.1.1 A site-specific survey measured pH levels at various depths in the water column across the Caledonia North OECC (Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)), results of which are displayed in Table 5-2. The pH values remained relatively stable throughout the water column and across the Caledonia North OECC. However, there were slight increases in pH observed in conjunction with the presence of thermoclines. Air-sea interaction with carbon dioxide ensures seawater is generally within a range of 7.5 to 8.5, which is in agreement with the survey output.

Table 5-2: Summary of pH throughout the water column across the Caledonia North OECC (Source: Volume 7B, Appendix 4-2: Environmental Baseline Report (Offshore Export Cable Corridor)).

| Sample Station | Range of pH |     |         | pH at Specific Depth |     |           |
|----------------|-------------|-----|---------|----------------------|-----|-----------|
|                | Min         | Max | Surface | 25m                  | 50m | Max Depth |
| ENV53          | 8.0         | 8.0 | 8.0     | N/A                  | N/A | 8.0       |
| ENV17          | 7.9         | 8.0 | 8.0     | 7.9                  | 7.9 | 7.9       |
| ENV26          | 7.9         | 8.0 | 8.0     | 8.0                  | 7.9 | 7.9       |
| ENV04          | 7.9         | 8.1 | 8.1     | 8.0                  | 7.9 | 7.9       |
| ENV13          | 7.9         | 8.1 | 8.0     | 8.0                  | 7.9 | 7.9       |
| ENV29          | 7.9         | 8.1 | 8.1     | 8.0                  | 7.9 | 7.9       |
| ENV31          | 7.9         | 8.1 | 8.1     | 8.0                  | 7.9 | 7.9       |
| ENV36          | 7.9         | 8.1 | 8.1     | 8.0                  | 7.9 | 7.9       |



## 6 References

- <sup>1</sup> Marine Directorate (2024) 'The Marine Scotland NMPi Maps'. Available at: <https://marinescotland.atkinsgeospatial.com/nmpi> (Accessed 21/05/2024)
- <sup>2</sup> Sharples, J. *et al.* (2006) 'Inter-annual variability in the timing of stratification and the spring bloom in the north-western North Sea'. *Continental Shelf Research* 26(6) 733–751
- <sup>3</sup> Mahaffey, C., Palmer, M., Greenwood, N. and Sharples, J. (2020) 'Impacts of climate change on dissolved oxygen concentration relevant to the coastal and marine environment around the UK'. *MCCIP Science Review* 2020: 31–53
- <sup>4</sup> OSPAR (2023) 'Quality Status Report'. Available at: <https://www.ospar.org/work-areas/cross-cutting-issues/qsr2023> (Accessed 16/07/2024)
- <sup>5</sup> OSPAR (2017) 'Intermediate Assessment 2017'. Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/> (Accessed 30/05/2024)
- <sup>6</sup> United Kingdom Monitoring and Assessment Strategy (2018) 'Turbidity - Marine online assessment tool'. Available at: <https://moat.cefas.co.uk/ocean-processes-and-climate/turbidity/> (Accessed 21/05/24)
- <sup>7</sup> Centre for Environment, Fisheries and Aquaculture Science (Cefas) (2016) 'Suspended Sediment Climatologies around the UK'. Available at: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/584621/CEFAS\\_2016\\_Suspended\\_Sediment\\_Climatologies\\_around\\_the\\_UK.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/584621/CEFAS_2016_Suspended_Sediment_Climatologies_around_the_UK.pdf) (Accessed 20/05/24)
- <sup>8</sup> Copernicus Marine Service (2024) 'Atlantic – European North West Shelf – Ocean Biogeochemistry Analysis and Forecast'

Caledonia Offshore Wind Farm  
5th Floor, Atria One  
144 Morrison Street  
Edinburgh  
EH3 8EX

[www.caledoniaoffshorewind.com](http://www.caledoniaoffshorewind.com)

