



Code: UKCAL-CWF-CON-EIA-RPT-00008-1010

Volume 8 Additional Information

Appendix 10: Ornithology Apportioning Technical Note (Caledonia OWF)

Caledonia Offshore Wind Farm Ltd

5th Floor Atria One, 144 Morrison Street, Edinburgh, EH3 8EX

Volume 8 Appendix 10: Ornithology Apportioning Technical Note (Caledonia OWF)

Code	UKCAL-CWF-CON-EIA-RPT-00008-1010
Revision	Issued
Date	30 September 2025

Table of Contents

1	Introduction	1
2	Methods.....	2
2.1	Overview	2
2.2	Species and Relevant Colonies.....	2
2.3	Definitions of Seasons	11
2.4	Apportioning Methods	11
2.4.1	Overview	11
2.4.2	Breeding Season	12
2.4.3	Non-breeding Season	15
3	Results	16
3.1	Breeding season	16
3.1.1	Overview	16
3.1.2	Kittiwake	16
3.1.3	Guillemot	16
3.1.4	Razorbill	20
3.1.5	Puffin	20
3.1.6	Gannet.....	20
3.2	Non-breeding Season.....	24
3.2.1	Overview	24
3.2.2	Kittiwake	24
3.2.3	Great Black-Backed Gull	24
3.2.4	Herring Gull	27
3.2.5	Guillemot	27
3.2.6	Razorbill	29
3.2.7	Puffin	30
3.2.8	Gannet.....	31
4	Conclusion	33
	References	34

List of Tables

Table 2-1: Species and impact pathways for which apportioning was undertaken.	2
Table 2-2: Species specific foraging ranges as per Woodward <i>et al.</i> (2019). ...	3
Table 2-3: Designated colonies (feature or assemblage) identified for apportioning based on breeding season foraging range and/or connectivity during the non-breeding season, where the around coast distance to designated colonies has been measured from the centre of the Caledonia OWF to the centre of the designated colony.	5
Table 2-4: Defined seasons in the Scottish Marine Environment used in the assessment for key species (NatureScot, 2020).	11
Table 2-5: Proportion of adult, immature and sabbatical birds included within the apportionment process.	14
Table 3-1: Kittiwake breeding season (Mid-April to August) apportioning weightings to designated sites.	17
Table 3-2: Guillemot breeding season (April to mid-August) apportioning weightings to designated sites.	19
Table 3-3: Razorbill breeding season (April to mid-August) apportioning weightings to designated sites.	21
Table 3-4 Puffin breeding season (April to Mid- August) apportioning weightings to designated sites.	22
Table 3-5: Gannet breeding season (Mid-March to September) apportioning weightings to designated sites.	23
Table 3-6: Kittiwake apportionment of adult mortality within the UK North Sea BDMPS during the non-breeding season (Autumn migration: September to December; Spring migration: January to Early-April).	25
Table 3-7: Great black-backed gull apportionment of adult mortality within the UK North Sea BDMPS during the non-breeding season (September to March).	26
Table 3-8: Herring gull apportionment of adult mortality within a regional population during the non-breeding season (September to March).	27
Table 3-9: Guillemot apportionment of adult mortality within a regional population during the non-breeding season (Mid-August to March).	28

Table 3-10: Razorbill apportionment of adult mortality within a regional population during the non-breeding season (Autumn/Spring migrations: Mid-August to October; January to March).	29
Table 3-11: Razorbill apportionment of adult mortality within a regional population during the non-breeding season (Winter: November to December).	30
Table 3-12: Puffin apportionment of adult mortality within a regional population during the non-breeding season (Mid-August to March).	31
Table 3-13. Gannet apportionment of adult mortality within a regional population during the non-breeding season (Autumn migration: October to November).	32
Table 3-14: Gannet apportionment of adult mortality within a regional population during the non-breeding season (Spring migration: December to Early-March).	32

Acronyms and Abbreviations

BTO	British Trust for Ornithology
BDMPS	Biologically Defined Minimum Population Scales
HRA	Habitats Regulations Appraisal
JNCC	Joint Nature Conservation Committee
MD-LOT	Marine Directorate – Licensing Operations Team
MMFR	Mean Max Foraging Range
MS-LOT	Marine Scotland Licensing Operations Team
MSS	Marine Scotland Science
OWF	Offshore Wind Farm
RIAA	Report to Inform Appropriate Assessment
SD	Standard Deviation
SMP	Seabird Monitoring Programme
SPA	Special Protection Area

1 Introduction

- 1.1.1.1 This appendix provides the updated information on the seasonal apportioning of seabirds for the array area of the Proposed Development (Offshore), referred to as the Caledonia Offshore Wind Farm (OWF) and forms part of the addendums submitted in response to the Marine Directorate – Licensing Operations Team (MD-LOT) request for Additional Information (dated 17 July 2025).
- 1.1.1.2 Background to the Proposed Development (Offshore) and the consent applications for Caledonia North and Caledonia South are presented within the original consent applications (Volume 1, Chapter 5: Proposed Development Phasing) and summarised within the covering addendum document (Volume 8: Caledonia Offshore Wind Farm EIAR and HRA Addendum).
- 1.1.1.3 This document has been prepared specifically to inform the consent applications for the Proposed Development (Offshore), namely Caledonia North and Caledonia South, and presents the apportioning of predicted impacts to designated breeding seabird colonies, specifically those associated with Special Protection Areas (SPAs) and Ramsar sites, as well as non-designated breeding colonies.
- 1.1.1.4 Apportioning is the process by which the predicted impacts of the Proposed Development (Offshore) are allocated in a justifiable way to individual colonies.

2 Methods

2.1 Overview

2.1.1.1 The NatureScot Guidance Note 10 regarding apportioning impacts to breeding seabird colonies is not currently available as part of the NatureScot guidance to support offshore wind applications (NatureScot, 2023¹). As such, the assessment presented within this Technical Note is primarily based on the NatureScot Interim Apportioning Guidance (NatureScot, 2018²) as advised by NatureScot within consultation workshops (07 March 2024 and 04 June 2025).

2.1.1.2 It is important to note that although the Proposed Development (Offshore) will be developed in two phases (Caledonia North and Caledonia South), the apportioning distances presented within this technical note are for the Caledonia OWF (i.e., the array area of the Proposed Development (Offshore)). As requested by NatureScot's representations, separate apportioning distances for Caledonia North and Caledonia South are presented in separate appendices (Volume 8, Appendices 11 and 12).

2.2 Species and Relevant Colonies

2.2.1.1 Table 2-1 presents the seabird species for which apportioning was required based on the screening conclusions of the Proposed Development (Offshore), alongside the relevant impact pathway assessed for each species.

Table 2-1: Species and impact pathways for which apportioning was undertaken.

Species	Scientific Names	Nature of Impact
Kittiwake	<i>Rissa tridactyla</i>	Collision and distributional responses
Great black-backed gull	<i>Larus marinus</i>	Collision
Herring gull	<i>Larus argentatus</i>	Collision
Guillemot	<i>Uria aalge</i>	Distributional responses
Razorbill	<i>Alca torda</i>	Distributional responses
Puffin	<i>Fratercula arctica</i>	Distributional responses
Gannet	<i>Morus bassanus</i>	Collision and distributional responses

2.2.1.2 The species identified in Table 2-1 are protected as features of breeding designated colonies. It is noted that all designated sites screened in for assessment are SPAs, as such no Ramsar Sites are included within this appendix. Each bird has been apportioned to the designated sites within species-specific breeding season foraging ranges of the Caledonia OWF, defined as the mean maximum foraging range (MMFR) plus 1 standard deviation (SD) presented in Woodward *et al.* (2019³) (Table 2-2). It should be noted that the Interim Guidance from NatureScot (2018²) calls for the use of MMFR presented in Thaxter *et al.* (2012⁴); however, this has been superseded by Woodward *et al.* (2019³).

Table 2-2: Species specific foraging ranges as per Woodward *et al.* (2019³).

Species	Mean Max (km)	1 SD (km)	MMFR + 1SD (km)
Kittiwake	156.1	144.5	300.6
Great black-backed gull	73.0	-	73.0
Herring gull	58.8	26.8	85.6
Guillemot	73.2 (55.5)*	80.5 (39.7)*	153.7 (95.2)*
Razorbill	88.7 (73.8)*	75.9 (48.4)*	164.6 (122.2)*
Puffin	137.1	128.3	265.4
Gannet**	315.2	194.2	509.4
<p>* Distances in brackets exclude data from Fair Isle where foraging range may have been unusually high as a result of reduced prey availability during the study year (Woodward <i>et al.</i>, 2019³). These foraging ranges were used for sites south of the Pentland Firth for guillemot and razorbill as per the NatureScot Guidance Note 3 (NatureScot, 2023¹).</p> <p>** In line with NatureScot Guidance Note 3 (NatureScot, 2023¹), site specific maximum foraging ranges were used to assess potential connectivity to Forth Islands SPA, Grassholm SPA and St Kilda SPA. Forth Islands SPA and St Kilda SPA were within site specific maximum foraging ranges. As such Forth Islands has been assessed for gannet; however, St Kilda SPA was screened out due to the lack of expected connectivity when considering tracking data and literature evidence (as outlined within Application Document 13: Caledonia North Report to Inform Appropriate Assessment and Application Document 14: Caledonia South Report to Inform Appropriate Assessment).</p>			

- 2.2.1.3 A list of designated sites within MMFR + 1SD of the Proposed Development (Offshore) were identified within the updated HRA Screening Report (Volume 8, Appendix 7). This list was used to identify the sites considered in the apportioning assessment presented in this appendix. Table 2-3 provides the full list of designated sites and qualifying interests considered for apportioning.
- 2.2.1.4 The colony counts for the majority of designated and non-designated sites within foraging range (see Table 2-2) were derived from the British Trust for Ornithology (BTO) Seabird Monitoring Programme (SMP) database. Colony counts for East Caithness Cliffs SPA were derived from Burnell *et al.* (2023⁵) as requested by NatureScot within a consultation workshop regarding NatureScot representations following submission (04 June 2025). Colony counts are recorded as individuals, apparently occupied nests or apparently occupied sites; all counts recorded as apparently occupied nests or apparently occupied sites were treated as equivalent to pairs and were doubled to give the count as individuals. For guillemot and razorbill, a correction factor of x 1.34 was applied (Harris *et al.*, 2015⁶) to SMP counts to give total adult breeding numbers. For puffin a correction factor of x 2.00 was applied to SMP counts (Hughes *et al.* 2019⁷; Mitchell *et al.* 2004⁸). Where sites are made up of multiple sub-sites, the most recent available counts were added together, in this case, a range of years is presented within Table 2-3 below.

Table 2-3: Designated colonies (feature or assemblage) identified for apportioning based on breeding season foraging range and/or connectivity during the non-breeding season, where the around coast distance to designated colonies has been measured from the centre of the Caledonia OWF to the centre of the designated colony.

Designated Site	Species	Foraging Range (km) (MMFR + 1SD)	Distance to Designated Site (km)	Count of Breeding Adults (individuals)	Year(s) of Count
East Caithness Cliffs SPA**	Kittiwake	300.6	53.9	48,958	2015 - 2021
	Great black-backed gull*	73.0	54.0	532	2015 - 2021
	Herring gull*	85.6	53.7	6,600	2015 - 2021
	Guillemot	95.2	53.8	199,966 (149,228)	2015 - 2021
	Razorbill	122.2	53.7	40,373 (30,129)	2015 - 2021
North Caithness Cliffs SPA	Kittiwake	300.6	56.2 – 113.2	18,608	2015 - 2024
	Guillemot	95.2	55.2 – 113.2	62,102 (46,345)	2015 - 2024
	Razorbill	122.2	55.2 – 113.2	12,329 (9,201)	2015 - 2024
	Puffin	265.4	56.2 – 114.9	6,766 (3,383)	2015 - 2024
Troup, Pennan and Lion's Heads SPA	Kittiwake	300.6	59.4 – 62.3	27,344	2017 - 2023
	Herring gull*	85.6	59.4 – 62.3	1,108	1995 - 2023
	Guillemot	95.2	59.4 – 61.8	47,719 (35,611)	2017 - 2023
	Razorbill	122.2	59.4 – 62.3	8,801 (6,568)	2017 - 2023

Designated Site	Species	Foraging Range (km) (MMFR + 1SD)	Distance to Designated Site (km)	Count of Breeding Adults (individuals)	Year(s) of Count
Copinsay SPA	Kittiwake	300.6	80.7	670	2023
	Guillemot	153.7	80.7 – 81.7	1,312 (979)	2015 - 2024
	Great black-backed gull*	73.0	80.7 – 81.7	97	2023 - 2024
Hoy SPA	Kittiwake	300.6	82.7 – 97.7	608	2016 – 2017
	Guillemot	153.7	82.7 – 100.6	16,345 (12,198)	2016 - 2017
	Puffin	265.4	82.7 – 100.6	722 (361)	2016 - 2017
	Great black-backed gull*	73.0	84.8 – 95.5	775	1996 - 2024
Buchan Ness to Collieston Coast SPA	Kittiwake	300.6	103.8	27,094	2023
	Guillemot	95.2	103.8	40,763 (30,420)	2023
	Herring gull*	85.6	103.8	4,536	2023
Rousay SPA	Kittiwake	300.6	120.1 – 125.6	962	2016 - 2021
	Guillemot	153.7	120.1 – 126.4	7,921 (5,911)	2016 - 2018
Marwick Head SPA	Kittiwake	300.6	117.5	2,878	2023
	Guillemot	153.7	117.5	12,800 (9,552)	2023

Designated Site	Species	Foraging Range (km) (MMFR + 1SD)	Distance to Designated Site (km)	Count of Breeding Adults (individuals)	Year(s) of Count
Calf of Eday SPA	Kittiwake	300.6	117.8 – 121.1	324	2018 - 2024
	Guillemot	153.7	117.8 – 122.2	7,402 (5,524)	2018
West Westray SPA	Kittiwake	300.6	131.1 – 135.1	8,004	1999 - 2023
	Guillemot	153.7	129.4 – 137.4	43,035 (32,116)	2017 - 2023
	Razorbill	164.6	129.4 – 137.4	3,103 (2,316)	2017 - 2023
Fowlsheugh SPA	Kittiwake	300.6	159.4 – 163.2	40,156	2018 - 2023
Cape Wrath SPA	Kittiwake	300.6	169.1 – 184.6	6,656	2000 - 2023
	Puffin	265.4	167.5 – 176.1	428 (214)	2017 - 2023
Sule Skerry and Sule Stack SPA	Puffin	265.4	154.3	95,484 (47,742)	2018
	Gannet	509.4	154.3 – 155.4	15,648	2013 - 2024
	Guillemot	95.2	154.3 – 155.5	14,284 (10,660)	1998 - 2024

Designated Site	Species	Foraging Range (km) (MMFR + 1SD)	Distance to Designated Site (km)	Count of Breeding Adults (individuals)	Year(s) of Count
Fair Isle SPA	Kittiwake	300.6	161.0	896	2021
	Razorbill	164.6	161.0	2,580 (1,925)	2021
	Puffin	265.4	161.0	13,332 (6,666)	2015
	Gannet	509.4	161.0	11,184	2024
	Guillemot	153.7	161.0	24,515 (18,295)	2021
Sumburgh Head SPA	Kittiwake	300.6	202.0 – 205.7	691	2018 – 2023
Foula SPA	Kittiwake	300.6	222.5	1,193	2021
	Puffin	265.4	222.5	12,702 (6,351)	2016
North Rona and Sula Sgeir SPA	Kittiwake	300.6	234.4 – 252.6	1,424	2021
	Puffin	265.4	234.4 – 252.6	5,668 (2,834)	2021
	Gannet	509.4	252.6	18,990	2023
Forth Islands SPA***	Kittiwake	300.6	248.5 – 264.6	14,216	2019 - 2024
	Gannet	509.4	261.7	162,000	2021
	Puffin	265.4	248.5 – 291.5	117,960 (58,980)	2024

Designated Site	Species	Foraging Range (km) (MMFR + 1SD)	Distance to Designated Site (km)	Count of Breeding Adults (individuals)	Year(s) of Count
Noss SPA	Kittiwake	300.6	239.6	154	2024
	Puffin	265.4	234.7 – 239.6	1,194 (597)	2000 - 2023
	Gannet	509.4	239.6	24,670	2023
St Abb's Head to Fast Castle SPA	Kittiwake	300.6	271.8 – 273.4	11,992	2024
Hermaness, Saxa Vord and Valla Field SPA	Kittiwake	300.6	318.3 – 321.0	378	1999 - 2023
	Gannet	509.4	319.6	39,606	2024
Handa SPA	Kittiwake	300.6	209.4	9,178	2023
Shiant Isles SPA	Kittiwake	300.6	289.3 – 292.9	2,318	2015 - 2023
Farne Islands SPA	Kittiwake*	300.6	301.0	5,790	2024
Flamborough and Filey Coast SPA	Gannet*	509.4	489.4	31,588	2024

Designated Site	Species	Foraging Range (km) (MMFR + 1SD)	Distance to Designated Site (km)	Count of Breeding Adults (individuals)	Year(s) of Count
<p>* Species assessed for the non-breeding season only, approach agreed with NatureScot via email on 07 August 2025.</p> <p>** Colony counts taken from Burnell <i>et al.</i>, 2023⁵, approach agreed with NatureScot within a consultation meeting on 07 August 2025.</p> <p>*** The Forth Islands SPA colony count takes into account the 2021 estimated Bass Rock drone count of 81,000 AOS (Harris <i>et al.</i>, 2023⁹; Wanless <i>et al.</i>, 2023¹⁰), approach agreed with NatureScot within a consultation meeting on 07 August 2025.</p> <p>Note, foraging ranges presented in “()” exclude data from Fair Isle where foraging range may have been unusually high (Woodward <i>et al.</i>, 2019³). These foraging ranges were used for sites south of the Pentland Firth for guillemot and razorbill. Counts for guillemot and razorbill apply Harris <i>et al.</i> (2015⁶⁶) correction factor x 1.34 for conversion to total adult breeding numbers. Counts for puffin apply a correction factor x 2.00 for conversion to total adult breeding numbers (Hughes <i>et al.</i> 2019⁷⁷; Mitchell <i>et al.</i> 2004⁸⁸). SMP counts are shown in “()”. The distance of each colony was measured as the around land distance from the geometric centre of the Caledonia OWF to the geometric centre of the colony. When undertaking apportioning, colonies with multiple sites were treated separately and the geometric centre of each site within the colony was used (as such, distances presented differ between species depending on which sites had associated colony counts).</p>					

2.3 Definitions of Seasons

- 2.3.1.1 The defined breeding seasons for all species were derived from the NatureScot (2020¹¹) guidance (see Table 2-4).
- 2.3.1.2 Apportioning outside the breeding season relies on information from Biologically Defined Minimum Population Scales (BDMPS) (Furness, 2015¹²). Furness (2015¹²) identifies autumn passage, spring passage and winter periods within the non-breeding season for kittiwake, gannet, and razorbill, as such apportioning weightings were derived for these periods using the period within which the mean peak was recorded for each species. The non-breeding defined seasons used within apportioning for kittiwake, razorbill, and gannet are presented in Table 2-4. The Furness (2015¹²) seasons were foreshortened to reduce overlap with the NatureScot breeding season definitions to avoid overestimating seasonal mortality estimates.

Table 2-4: Defined seasons in the Scottish Marine Environment used in the assessment for key species (NatureScot, 2020¹¹).

Species	Breeding Season	Non-breeding Season
Kittiwake	Mid-April to August	September to Early April
Great black-backed gull	April to August	September to March
Herring gull	April to August	September to March
Common guillemot	April to mid-August	Late August to March
Razorbill	April to mid-August	Late August to March
Puffin	April to Mid- August	Late August to March
Gannet	Mid-March to September	October to Early-March

2.4 Apportioning Methods

2.4.1 Overview

- 2.4.1.1 The weightings obtained from the apportioning were used to allocate seasonal mortality estimates for each species by age class to seabird colonies.

2.4.2 Breeding Season

- 2.4.2.1 The assessment presented within this Technical Note is primarily based on the NatureScot Interim Guidance (NatureScot, 2018²) regarding apportioning impacts to breeding seabird colonies.
- 2.4.2.2 During the breeding season there is potential not only for breeding adult birds within foraging range of the project to have connectivity but also juvenile, immature and sabbatical birds which are not associated with any given colony. The first step in the apportionment process is therefore to calculate the proportion of breeding adult birds for which the project may impact.
- 2.4.2.3 The proportion of juvenile and immature birds comparatively to the number of breeding adults which may be connected to the project can typically be calculated using the age ratios from the digital aerial surveys or from using generalised stable age structure data. In relation to the use of age ratios from the digital aerial survey data, there are a number of key issues with accurately identifying age of key seabirds as detailed below.
- 2.4.2.4 For kittiwakes only first winter juvenile birds are readily distinguishable from other age categories, due to the distinct 'W pattern' across the wings and black tail-band (Svensson *et al.*, 2009¹³). This pattern, however, is lost by the time a kittiwake reaches its second winter moult, whereby the bird is indistinguishable from an adult bird. As presented in Coulson (2011¹⁴), the modal age of kittiwakes first breeding is four years old, although the age of first breeding has been documented as late as 10 years old. This clearly shows that by simply applying the assumption that all adult plumage birds are breeding adults, as would be the case when using site-specific survey data, it is highly likely to overestimate the proportion of breeding adult birds within the Caledonia OWF.
- 2.4.2.5 For guillemot, razorbill and puffin only first winter juvenile birds are readily distinguishable from other age categories, with the distinguishing feature of juveniles being their size in comparison to the adult males in attendance of the chicks. After their first winter immature auks are indistinguishable from breeding adult birds. The average first breeding age for guillemot, razorbill and puffin is six, five and five years old respectively (Horswill and Robinson, 2015¹⁵). Therefore, the treatment that all 'adult type' appearance birds are breeding adults, as would be the case when using site-specific survey data, it is highly likely to overestimate the proportion of breeding adult birds with the project area.
- 2.4.2.6 In relation to gannet, with juvenile (first calendar year birds) plumage being primarily grey/brown in colour with a lack of a distinct yellow head (Svensson *et al.*, 2009¹³) this makes them distinctly different to adult birds. For second calendar year birds, the grey-brown plumage on the head, underparts, uppertail-coverts and usually some of the lesser wing uppertail-coverts becomes white (Svensson *et al.*, 2009¹³), makes this age

category readily distinguishable from adult birds. For third calendar year birds most tail-feathers and secondaries are usually black intermixed with white feathers, whilst the remaining body and head largely resemble the plumage of an adult bird, although these birds are still readily identifiable from adult birds, depending on the quality of the survey data and behaviour of the bird recorded (e.g., banking birds) might be difficult to observe and therefore this age category may be less regularly distinguished from adult birds. For fourth calendar year birds only the central tail-feathers and the odd scattered secondaries remain black, the rest of the bird's plumage resembles that of an adult bird, depending on the quality of the survey data and behaviour of the bird recorded (e.g., banking birds) might be difficult to observe and therefore this age category may be less regularly distinguished from adult birds. From fourth calendar year onwards the plumage of gannets remains indistinguishable, with the average age of first breeding at five years old. There is therefore potential to overestimate the proportion of breeding adult birds with the project area when using site-specific survey data.

- 2.4.2.7 Furthermore, not all adult birds within the Caledonia OWF can be classified as breeding birds. This is evidenced from adult sabbatical birds free roaming the North Sea whilst taking a break from breeding activities (Marine Scotland, 2017¹⁶). A sabbatical rate of 10% for gannet and kittiwake populations and 7% for auk species was advocated by Marine Scotland based on expert opinion for inclusion within other Scottish projects such as Moray West and other Forth and Tay projects and therefore has been applied to assessments for the Proposed Development (Offshore) also. Sabbatical rates for great black-backed gull and herring gull were derived from Horswill and Robinson (2015¹⁵).
- 2.4.2.8 The data presented in Furness (2015¹²) are considered to provide a more accurate representation of population age structure than site-specific survey data for reasons set out above. Furthermore, Furness (2015¹²) draws upon a wide number of data sources gathered across multiple years to model population age structure, thus reducing the potential for any bias associated with the snapshot nature of site-based surveys. A summary of the adult/immature age ratio for key seabirds is provided in Table 2-5.

Table 2-5: Proportion of adult, immature and sabbatical birds included within the apportionment process.

Species	Adult/Immature Ratio	Sabbatical Rate	Total Breeding Adult Percentage
Kittiwake	53% / 47%	10%	47.70%
Guillemot	57% / 43%	7%	53.01%
Razorbill	57% / 43%	7%	53.01%
Puffin	55% / 45%	7%	51.15%
Gannet	55% / 45%	10%	49.50%

- 2.4.2.9 Once the number of breeding adults potentially impacted has been calculated, the next step is to apportion impacts from the Caledonia OWF to each specific breeding colony with potential connectivity. This was undertaken following NatureScot's (2018²) interim guidance. The methodology calculates a proportional weighting for each colony based on the following three weighting factors:
- The population size of each colony (individuals);
 - The distance from the center of the proposed array area to the center of the colony; and
 - The available sea area within MMFR + 1SD of the colony, expressed as a proportion of total potential area.

- 2.4.2.10 These weighting factors are combined to form the following equation:

$$Weight = \left(\frac{Colony\ Population}{Sum\ of\ Populations} \right) \times \left(\frac{Sum\ of\ Distance^2}{Colony\ Distance^2} \right) \times \left(\frac{\frac{1}{Colony\ Sea\ Proportion}}{Sum\ of\ \frac{1}{Sea\ Proportions}} \right)$$

- 2.4.2.11 The population estimates at the relevant designated sites included in the calculation are detailed Table 2-3. Population estimates at additional colonies and non-designated sites are presented within Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF). These counts were obtained from the BTO SMP database, with the exception of counts made of East Caithness Cliffs SPA. For the East Caithness Cliffs SPA, counts from were derived from Burnell *et al.* (2023⁵) as requested by NatureScot within a consultation workshop regarding NatureScot representations following submission (04 June 2025).
- 2.4.2.12 Foraging ranges are based on at sea distances taking into account land barriers to movements for species which are known to avoid commuting over land.

- 2.4.2.13 Distances between colonies and the Caledonia OWF, and the percentage of area at sea for each colony, were calculated in QGIS. As per the NatureScot (2018²) Interim Guidance, the distance of the colony was measured as the around land distance from the geometric centre of the Caledonia OWF to the geometric centre of the colony, taking the shortest at sea distance route possible to identify colonies for inclusion within assessments. When undertaking apportioning, colonies with multiple sites were treated separately and the geometric centre of each site within the colony was used. Sea area was calculated by buffering the colony centroid by the seabirds MMFR + 1SD then removing all area over land and areas where seabirds are unlikely to forage such as estuaries.

2.4.3 Non-breeding Season

- 2.4.3.1 The BDMPS population estimates were used to apportion all birds present at the Caledonia OWF during the non-breeding season to breeding colony populations for all species, except for guillemot (Furness, 2015¹²). The BDMPS and associated abundance estimates presented in Furness (2015¹²¹²¹²¹²) are based upon data on demography, migration and modelled population age structure from the UK and overseas. For most species, the relevant BDMPS region for the Caledonia OWF is the UK North Sea and Channel waters, except for kittiwake and great black-backed gull for which the UK North Sea waters is the most relevant BDMPS region.
- 2.4.3.2 For guillemot, apportioning for the non-breeding season was based on the breeding population found within the MMFR + 1SD of the Caledonia OWF. This is in line with the approach outlined in the NatureScot Guidance Note 3 (NatureScot, 2023¹), based on recent geolocator studies presented in Buckingham *et al.* (2022¹⁷).

3 Results

3.1 Breeding season

3.1.1 Overview

3.1.1.1 The apportioning weightings of predicted impacts from the Caledonia OWF for all designated site qualifying features screened in for assessment are provided within this section for the breeding season (Table 3-1 to Table 3-5). Full lists of colonies and apportioning weightings of predicted impacts for non-designated sites are provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).

3.1.2 Kittiwake

3.1.2.1 The weightings for apportioning breeding season kittiwake to designated sites within MMFR+1SD of the Caledonia OWF are provided in Table 3-1. These weightings were used to apportion the potential number of kittiwake distributional response and collision mortalities during the breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

3.1.3 Guillemot

3.1.3.1 The weightings for apportioning breeding season guillemot to designated sites within MMFR+1SD of the Caledonia OWF are provided in Table 3-2. These weightings were used to apportion the potential number of guillemot distributional response mortalities during the breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-1: Kittiwake breeding season (Mid-April to August) apportioning weightings to designated sites.

Site Name	SMP Database Counts (Individuals)	Distance to Caledonia OWF (km)	1/Proportion of Foraging Range at Sea	Resulting Weight	Proportional Weight
East Caithness Cliffs SPA*	48,958	53.9	0.02	3.039	0.483
North Caithness Cliffs SPA	18,608	55.2 – 113.2	0.20	0.395	0.063
Troup, Pennan and Lion's Heads SPA	27,344	59.4 – 62.3	0.09	1.340	0.213
Copinsay SPA	670	80.7	0.01	0.014	0.002
Hoy SPA	608	82.7 – 97.7	0.09	0.012	0.002
Buchan Ness to Collieston Coast SPA	27,094	103.8	0.02	0.445	0.071
Rousay SPA	962	120.1 – 125.6	0.11	0.009	0.001
Marwick Head SPA	2,878	117.5	0.01	0.029	0.005
Calf of Eday SPA	324	117.8 – 121.1	0.05	0.003	<0.001
West Westray SPA	8,004	131.1 – 135.1	0.06	0.061	0.010
Fowlsheugh SPA	40,156	159.4 – 163.2	0.07	0.327	0.052
Cape Wrath SPA	6,656	169.1 – 184.6	0.35	0.034	0.005
Fair Isle SPA	896	161.0	0.01	0.004	0.001
Sumburgh Head SPA	691	202.0 – 205.7	0.04	0.002	<0.001
Foula SPA	1,193	222.5	0.01	0.003	<0.001
North Rona and Sula Sgeir SPA	1,424	234.5 – 252.6	0.02	0.003	0.001

Site Name	SMP Database Counts (Individuals)	Distance to Caledonia OWF (km)	1/Proportion of Foraging Range at Sea	Resulting Weight	Proportional Weight
Forth Islands SPA	14,216	248.5 – 264.6	0.15	0.062	0.010
Noss SPA	154	239.6	0.01	<0.001	<0.001
St Abb's Head to Fast Castle SPA	11,992	271.8 – 273.4	0.09	0.041	0.006
Handa SPA	9,178	209.4	0.01	0.033	0.005
Shiant Isles SPA	2,318	289.3 – 292.9	0.07	0.005	0.001
Hermaness, Saxa Vord and Valla Field SPA	378	318.3 - 321.0	0.05	<0.001	<0.001
Additional colonies and non-designated sites	50,680	45.9 – 347.8	3.66	0.434	0.069
<p>Note the residual weight assigned to non-SPA colonies are presented here as an overall total. Further breakdown of these sites is provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).</p> <p>* Colony counts taken from Burnell <i>et al.</i> (2023⁵).</p>					

Table 3-2: Guillemot breeding season (April to mid-August) apportioning weightings to designated sites.

Site Name	SMP Database Counts (Individuals)	Breeding Individuals*	Distance to Caledonia OWF (km)	1/Proportion of Foraging Range at Sea	Resulting Weight	Proportional Weight
East Caithness Cliffs SPA**	149,228	199,966	53.8	0.02	2.178	0.707
North Caithness Cliffs SPA	46,345	62,102	55.2 – 113.2	0.27	0.286	0.093
Troup, Pennan and Lion's Heads SPA	35,611	47,719	59.4 – 61.8	0.08	0.327	0.106
Copinsay SPA	979	1,312	80.7 – 81.7	0.03	0.004	0.001
Hoy SPA	12,198	16,345	82.7 – 100.6	0.14	0.041	0.013
Buchan Ness to Collieston Coast SPA	30,420	40,763	103.8	0.01	0.089	0.029
Rousay SPA	5,911	7,921	120.1 – 126.4	0.11	0.009	0.003
Marwick Head SPA	9,552	12,800	117.5	0.01	0.017	0.006
Calf of Eday SPA	5,524	7,402	117.8 – 122.2	0.08	0.009	0.003
West Westray SPA	32,116	43,035	129.4 - 137.4	0.07	0.043	0.014
Sule Skerry and Sule Stack SPA	10,660	14,284	154.3 - 155.5	0.02	0.011	0.004
Fair Isle SPA	18,295	24,515	161.0	0.01	0.016	0.005
Additional colonies and non-designated sites	19,965	26,753	44.53– 138.0	0.60	0.062	0.020
<p>Note, the residual weight assigned to non-SPA colonies are presented here as an overall total. Further breakdown of these sites is provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).</p> <p>* Applying Harris <i>et al.</i> (2015⁶) correction factor x 1.34 for total adult breeding numbers.</p> <p>** Colony counts taken from Burnell <i>et al.</i> (2023⁵)</p>						

3.1.4 Razorbill

- 3.1.4.1 The weightings for apportioning breeding season razorbill to designated sites within MMFR+1SD of the Caledonia OWF are provided in Table 3-3. These weightings were used to apportion the potential number of razorbill distributional response mortalities during the breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

3.1.5 Puffin

- 3.1.5.1 The weightings for apportioning breeding season puffin to designated sites within MMFR+1SD of the Caledonia OWF are provided in Table 3-4. These weightings were used to apportion the potential number of puffin distributional response mortalities during the breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

3.1.6 Gannet

- 3.1.6.1 The weightings for apportioning breeding season gannet to designated sites within MMFR+1SD of the Caledonia OWF are given in Table 3-5. These weightings were used to apportion the potential number of gannet distributional response and collision mortalities to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-3: Razorbill breeding season (April to mid-August) apportioning weightings to designated sites.

Site Name	SMP Database Counts (Individuals)	Breeding Individuals*	Distance to Caledonia OWF (km)	1/Proportion of Foraging Range at Sea	Resulting Weight	Proportional Weight
East Caithness Cliffs SPA**	30,129	40,373	53.7	0.02	2.340	0.724
North Caithness Cliffs SPA	9,201	12,329	55.2 – 113.2	0.28	0.267	0.081
Troup, Pennan and Lion's Heads SPA	6,568	8,801	59.4 – 62.3	0.10	0.327	0.099
West Westray SPA	2,316	3,103	129.4 – 137.4	0.08	0.016	0.005
Fair Isle SPA	1,925	2,580	161.0	0.01	0.009	0.003
Additional colonies and non-designated sites	16,835	22,559	45.9 – 144.0	1.85	0.290	0.088
<p>Note the residual weight assigned to non-SPA colonies are presented here as an overall total. Further breakdown of these sites is provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).</p> <p>* Applying Harris <i>et al.</i> (2015⁹) correction factor x 1.34 for total adult breeding numbers.</p> <p>** Colony counts taken from Burnell <i>et al.</i> (2023⁵).</p>						

Table 3-4 Puffin breeding season (April to Mid- August) apportioning weightings to designated sites.

Site Name	SMP Database Counts (Individuals)	Breeding Individuals*	Distance to Caledonia OWF (km)	1/Proportion of Foraging Range at Sea	Resulting Weight	Proportional Weight
North Caithness Cliffs SPA	3,383	6,766	55.2 – 114.9	0.27	0.076	0.045
Hoy SPA	361	722	82.7 – 100.6	0.10	0.014	0.008
Cape Wrath SPA	214	428	167.5 – 176.1	0.35	0.003	0.002
Sule Skerry and Sule Stack SPA	47,742	95,484	154.3 - 155.4	0.01	0.360	0.213
Fair Isle SPA	6,666	13,332	161.0	0.01	0.082	0.048
Foula SPA	6,351	12,702	222.5	0.01	0.039	0.023
North Rona and Sula Sgeir SPA	2,834	5,668	234.4 – 252.6	0.02	0.009	0.005
Forth Islands SPA	58,980	117,960	248.5 - 291.5	0.17	0.673	0.398
Noss SPA	597	1,194	234.7 - 239.6	0.02	0.002	0.001
Additional colonies and non-designated sites	76,257	152,514	45.9 – 445.8	3.64	0.435	0.257
<p>Note, the residual weight assigned to non-SPA colonies are presented here as an overall total. Further breakdown of these sites is provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).</p> <p>* Applying correction factor x 2.00 for total adult breeding numbers (Hughes <i>et al.</i> 2019⁷; Mitchell <i>et al.</i> 2004⁸⁸).</p>						

Table 3-5: Gannet breeding season (Mid-March to September) apportioning weightings to designated sites.

Site Name	SMP Database Counts (Individuals)	Distance to Caledonia OWF (km)	1/Proportion of Foraging Range at Sea	Resulting Weight	Proportional Weight
Sule Skerry and Sule Stack SPA	15,648	154.3 - 155.4	0.02	0.099	0.059
Fair Isle SPA	11,184	161.0	0.01	0.068	0.040
North Rona and Sula Sgeir SPA	18,990	252.6	0.01	0.045	0.027
Forth Islands SPA*	162,000	261.7	0.02	0.658	0.391
Noss SPA	24,670	239.6	0.01	0.068	0.041
Hermaness, Saxa Vord and Valla Field SPA	39,606	319.6	0.01	0.060	0.036
Additional colonies and non-designated sites	162,919	59.6 - 489.4	0.14	0.682	0.406
<p>Note, the residual weight assigned to non-SPA colonies are presented here as an overall total. Further breakdown of these sites is provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).</p> <p>* The Forth Islands SPA colony count takes into account the 2021 estimated Bass Rock drone count of 81,000 AOS (Harris <i>et al.</i>, 2023⁹; Wanless <i>et al.</i>, 2023¹⁰), approach agreed with NatureScot within a consultation meeting on 07 August 2025.</p>					

3.2 Non-breeding Season

3.2.1 Overview

- 3.2.1.1 The apportioning weightings of predicted impacts from the Caledonia OWF for all designated site qualifying features screened in for assessment are provided within this section for the non-breeding season (Table 3-6 to Table 3-14).

3.2.2 Kittiwake

- 3.2.2.1 The non-breeding defined seasons for kittiwake were based on Furness (2015¹²), which were modified to align with the NatureScot (2020¹¹) breeding season periods (Table 2-4).
- 3.2.2.2 The weightings for apportioning non-breeding season kittiwake to designated sites are provided in Table 3-6 for autumn migration (September to December), and for spring migration (January to Early-April) and were based on the UK North Sea BDMPS abundance data. These weightings were used to apportion the potential numbers of non-breeding kittiwake distributional response and collision mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

3.2.3 Great Black-Backed Gull

- 3.2.3.1 The non-breeding defined seasons for great black-backed gull were based on NatureScot (2020¹¹) guidance (Table 2-4).
- 3.2.3.2 The weightings for apportioning non-breeding season great black-backed gull to designated sites are provided in Table 3-7 for non-breeding season (September to March) and were based on the UK North Sea BDMPS abundance data. These weightings were used to apportion the potential numbers of non-breeding great black-backed gull collision mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-6: Kittiwake apportionment of adult mortality within the UK North Sea BDMPs during the non-breeding season (Autumn migration: September to December; Spring migration: January to Early-April).

Site Name	UK North Sea Adults Associated with the SPA	Autumn Migration		Spring Migration	
		UK North Sea Total Birds	Proportional Weighting	UK North Sea Total Birds	Proportional Weighting
East Caithness Cliffs SPA	48,492	829,937	0.058	627,816	0.077
North Caithness Cliffs SPA	12,180	829,937	0.015	627,816	0.019
Troup, Pennan and Lion's Heads SPA	17,875	829,937	0.022	627,816	0.028
Copinsay SPA	799	829,937	0.001	627,816	0.001
Hoy SPA	476	829,937	0.001	627,816	0.001
Buchan Ness to Collieston Coast SPA	15,050	829,937	0.018	627,816	0.024
Rousay SPA	2,117	829,937	0.003	627,816	0.003
Marwick Head SPA	631	829,937	0.001	627,816	0.001
Calf of Eday SPA	896	829,937	0.001	627,816	0.001
West Westray SPA	14,466	829,937	0.017	627,816	0.023
Fowlsheugh SPA	11,204	829,937	0.013	627,816	0.018
Cape Wrath SPA	207	829,937	<0.001	627,816	<0.001
Fair Isle SPA	925	829,937	0.001	627,816	0.001
Sumburgh Head SPA	252	829,937	<0.001	627,816	<0.001
Foula SPA	392	829,937	<0.001	627,816	0.001

Site Name	UK North Sea Adults Associated with the SPA	Autumn Migration		Spring Migration	
		UK North Sea Total Birds	Proportional Weighting	UK North Sea Total Birds	Proportional Weighting
North Rona and Sula Sgeir SPA	25	829,937	<0.001	627,816	<0.001
Forth Islands SPA	3,720	829,937	0.004	627,816	0.006
Noss SPA	608	829,937	0.001	627,816	0.001
St Abb's Head to Fast Castle SPA	4,084	829,937	0.005	627,816	0.007
Handa SPA	37	829,937	<0.001	627,816	<0.001
Shiant Isles SPA	11	829,937	<0.001	627,816	0.000
Farne Islands SPA	4,123	829,937	0.005	627,816	0.007
Hermaness, Saxa Vord and Valla Field SPA	469	829,937	0.001	627,816	0.001

Table 3-7: Great black-backed gull apportionment of adult mortality within the UK North Sea BDMPS during the non-breeding season (September to March).

Site Name	UK North Sea Adults Associated with the SPA	Non-breeding Season	
		UK North Sea Total Birds	Proportional Weighting
Copinsay SPA	436	91,399	0.005
East Caithness Cliffs SPA	350	91,399	0.004
Hoy SPA	120	91,399	0.001

3.2.4 Herring Gull

- 3.2.4.1 The non-breeding defined seasons for herring gull were based on NatureScot (2020¹¹) guidance (Table 2-4).
- 3.2.4.2 The weightings for apportioning non-breeding season herring gull to designated sites are provided in Table 3-8 for non-breeding season (September to March), and were based on the UK North Sea and Channel waters BDMPs abundance data. These weightings were used to apportion the potential numbers of non-breeding herring gull collision mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-8: Herring gull apportionment of adult mortality within a regional population during the non-breeding season (September to March).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Non-breeding Season	
		UK North Sea Total Birds	Proportional Weighting
East Caithness Cliffs SPA	6,718	466,511	0.014
Troup, Pennan and Lion's Heads SPA	3,162	466,511	0.007
Buchan Ness to Collieston Coast SPA	6,166	466,511	0.013

3.2.5 Guillemot

- 3.2.5.1 The non-breeding defined seasons for guillemot were based on NatureScot (2020¹¹) guidance (Table 2-4).
- 3.2.5.2 The weightings for apportioning non-breeding season guillemot to designated sites are provided in Table 3-9 for non-breeding season (Mid-August to March) and were based on the UK North Sea and Channel waters BDMPs abundance data. These weightings were used to apportion the potential numbers of non-breeding guillemot distributional response mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-9: Guillemot apportionment of adult mortality within a regional population during the non-breeding season (Mid-August to March).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Non-breeding Season	
		UK North Sea Total Birds	Proportional Weighting
East Caithness Cliffs SPA	149,100	1,617,306	0.092
North Caithness Cliffs SPA	65,800	1,617,306	0.041
Troup, Pennan and Lion's Heads SPA	15,313	1,617,306	0.009
Copinsay SPA	7,850	1,617,306	0.005
Hoy SPA	8,820	1,617,306	0.005
Buchan Ness to Collieston Coast	20,658	1,617,306	0.013
Rousay SPA	8,680	1,617,306	0.005
Marwick Head SPA	15,536	1,617,306	0.010
Calf of Eday SPA	8,820	1,617,306	0.005
West Westray SPA	47,460	1,617,306	0.029
Sule Skerry and Sule Stack SPA	763	1,617,306	<0.001
Fair Isle SPA	18,929	1,617,306	0.012

3.2.6 Razorbill

- 3.2.6.1 The non-breeding defined seasons for razorbill were based on Furness (2015¹²), which were modified to align with the NatureScot (2020¹¹) breeding season periods (Table 2-4).
- 3.2.6.2 The weightings for apportioning non-breeding season razorbill to designated sites are provided Table 3-10 autumn/spring migrations (Mid-August to October; January to March), and Table 3-11 for winter (November to December), and were based on the UK North Sea and Channel waters BDMPS abundance data. These weightings were used to apportion the potential numbers of non-breeding razorbill distributional response mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-10: Razorbill apportionment of adult mortality within a regional population during the non-breeding season (Autumn/Spring migrations: Mid-August to October; January to March).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Autumn/Spring Migration	
		UK North Sea Total Birds	Proportional Weighting
East Caithness Cliffs SPA	25,000	591,874	0.042
North Caithness Cliffs SPA	3,230	591,874	0.005
Troup, Pennan and Lion's Heads SPA	3,486	591,874	0.006
West Westray SPA	1,045	591,874	0.002
Fair Isle SPA	1,738	591,874	0.003

Table 3-11: Razorbill apportionment of adult mortality within a regional population during the non-breeding season (Winter: November to December).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Winter	
		UK North Sea Total Birds	Proportional Weighting
East Caithness Cliffs SPA	7,500	218,622	0.034
North Caithness Cliffs SPA	1,020	218,622	0.005
Troup, Pennan and Lion's Heads SPA	1,046	218,622	0.005
West Westray SPA	330	218,622	0.002
Fair Isle SPA	549	218,622	0.003

3.2.7 Puffin

- 3.2.7.1 The non-breeding defined seasons for puffin were based on NatureScot (2020¹¹) guidance (Table 2-4).
- 3.2.7.2 The weightings for apportioning non-breeding season puffin to designated sites are provided in Table 3-12 for non-breeding season (Mid-August to March) and were based on the UK North Sea and Channel waters BDMPS abundance data. These weightings were used to apportion the potential numbers of non-breeding puffin distributional response mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-12: Puffin apportionment of adult mortality within a regional population during the non-breeding season (Mid-August to March).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Non-breeding Season	
		UK North Sea Total Birds	Proportional Weighting
North Caithness Cliffs SPA	293	231,957	0.001
Hoy SPA	1,050	231,957	0.005
Cape Wrath SPA	3	231,957	<0.001
Sule Skerry and Sule Stack SPA	119	231,957	0.001
Fair Isle SPA	3,212	231,957	0.014
Foula SPA	6,750	231,957	0.029
North Rona and Sula Sgeir SPA	11	231,957	<0.001
Forth Islands SPA	62,231	231,957	0.268
Noss SPA	241	231,957	0.001

3.2.8 Gannet

- 3.2.8.1 The non-breeding defined seasons for gannet were based on Furness (2015¹²), which were modified to align with the NatureScot (2020¹¹) breeding season periods (Table 2-4).
- 3.2.8.2 The weightings for apportioning non-breeding season gannet to designated sites are provided Table 3-13 autumn migration (October to November) and Table 3-14 for spring migration (December to Early-March), and were based on the UK North Sea and Channel waters BDMPS abundance data. These weightings were used to apportion the potential numbers of non-breeding gannet distributional response and collision mortalities during the non-breeding season to each of the designated sites (apportioned mortalities are presented within Volume 8, Appendix 4).

Table 3-13. Gannet apportionment of adult mortality within a regional population during the non-breeding season (Autumn migration: October to November).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Autumn Migration	
		UK North Sea Total Birds	Proportional Weighting
Sule Skerry and Sule Stack SPA	935	456,298	0.002
Fair Isle SPA	6,278	456,298	0.014
North Rona and Sula Sgeir SPA	1,845	456,298	0.004
Forth Islands SPA	110,964	456,298	0.243
Noss SPA	15,627	456,298	0.034
Hermaness, Saxa Vord and Valla Field SPA	38,965	456,298	0.085
Flamborough and Filey Coast SPA	22,122	456,298	0.048

Table 3-14: Gannet apportionment of adult mortality within a regional population during the non-breeding season (Spring migration: December to Early-March).

Site Name	UK North Sea and Channel Waters Adults Associated with the SPA	Spring Migration	
		UK North Sea Total Birds	Proportional Weighting
Sule Skerry and Sule Stack SPA	0	248,385	0.000
Fair Isle SPA	5,494	248,385	0.022
North Rona and Sula Sgeir SPA	0	248,385	0.000
Forth Islands SPA	77,675	248,385	0.313
Noss SPA	13,674	248,385	0.055
Hermaness, Saxa Vord and Valla Field SPA	34,094	248,385	0.137
Flamborough and Filey Coast SPA	15,485	248,385	0.062

4 Conclusion

- 4.1.1.1 This appendix provides details regarding the approach by which potential collision and distributional response mortalities were apportioned to designated and non-designated colonies to inform the updated RIAA conclusions for the Proposed Development (Offshore) (Volume 8, Appendix 4).
- 4.1.1.2 Apportioning has been undertaken using NatureScot (2018²) guidance for the following species:
- Kittiwake;
 - Herring gull;
 - Guillemot;
 - Razorbill;
 - Puffin; and
 - Gannet.
- 4.1.1.3 The weightings which have been used to apportion predicted impacts from the Caledonia OWF for all qualifying features screened in for assessment are provided within this Section 3.1 for the breeding season (Table 3-1 to Table 3-5) and Section 3.2 for the non-breeding season (Table 3-6 to Table 3-14). Full lists of colonies and non-designated sites (during the breeding season) are provided in Volume 8, Appendix 13: Ornithology Apportioning Results (Caledonia OWF).

References

- ¹ NatureScot (2023) 'Guidance Note 3: Guidance to support Offshore Wind applications: Marine Birds - Identifying theoretical connectivity with breeding site Special Protection Areas using breeding season foraging ranges'. Available at: <https://www.nature.scot/doc/guidance-note-3-guidance-support-offshore-wind-applications-marine-birds-identifying-theoretical> (Accessed 01/09/2024)
- ² NatureScot (2018) 'Interim Guidance on apportioning impacts from marine renewable developments to breeding seabird populations in SPAs'. Available at: <https://www.nature.scot/doc/interim-guidance-apportioning-impacts-marine-renewable-developments-breeding-seabird-populations> (Accessed 01/09/2024)
- ³ Woodward, I., Thaxter, C.B., Owen, E. and Cook, A.S.C.P. (2019) 'Desk-based revision of seabird foraging ranges used for HRA screening'. BTO research report number 724
- ⁴ Thaxter, C.B., Lascelles, B., Sugar, K., Cook, A.S., Roos, S., Bolton, M., Langston, R.H. and Burton, N.H. (2012) 'Seabird foraging ranges as a preliminary tool for identifying candidate Marine Protected Areas'. Biological Conservation 156: 53-61
- ⁵ Burnell, D., Perkins, A.J., Newton, S.F., Bolton, M., Tierney, T.D. and Dunn, T.E. (2023). 'Seabirds Count: a census of breeding seabirds in Britain and Ireland (2015-2021)'. Lynx Nature Books.
- ⁶ Harris, M. P., Heubeck, M., Newell, M. A. and Wanless, S. (2015) 'The need for year-specific correction factors (k values) when converting counts of individual Common Guillemots *Uria aalge* to breeding pairs'. Bird Study 62(2): 276-279
- ⁷ Hughes, R.D., Le Bouard, F., Bradbury, G. and Owen, E. (2019). 'A Census of the Atlantic Puffins *Fratercula arctica* breeding on Orkney in 2016'. Seabird 31: 56-63.
- ⁸ Mitchell, P. I., Newton, S. F., Ratcliffe, N. and Dunn, T. E. (2004). 'Seabird Populations of Britain and Ireland'. T and A D Poyser; London.
- ⁹ Harris, M.P., Burton, E., Lewis, S., Tyndall, A., Nichol, C.J., Wade, T. and Wanless, S. (2023) 'Count of Northern Gannets on the Bass Rock in June 2023'
- ¹⁰ Wanless, S. Harris, M.P. and Murray, S. (2023) 'Northern Gannet *Morus bassanus*'. In: Burnell, D., Perkins, A.J., Newton, S.F., Bolton, M, Tierney, T.D. & Dunn, T.D. (eds). Seabirds Count, A census of breeding seabirds in Britain and Ireland (2015–2021). Lynx, Barcelona
- ¹¹ NatureScot (2020) 'Guidance Note 9: Seasonal definitions for birds in the Scottish marine environment'. Available at: <https://www.nature.scot/doc/guidance-note-9-guidance-support-offshore-wind-applications-seasonal-periods-birds-scottish-marine> (Accessed 14/08/2025).

¹² Furness, R.W. (2015) 'Non-breeding season populations of seabirds in UK waters: Population sizes for Biologically Defined Minimum Population Scales (BDMPS)'. Natural England Commissioned Reports, Number 164

¹³ Svensson, L., Mullarney, K. and Zetterström, D. (2009) 'Collins Bird Guide. The Most Complete Guide to The Birds Of Britain And Europe'. 2nd Edition

¹⁴ Coulson, J., 2011. 'The kittiwake'. A&C Black

¹⁵ Horswill, C. and Robinson, R.A. (2015) 'Review of seabird demographic rates and density dependence'. JNCC Report No: 552, JNCC, Peterborough, ISSN 0963-8901

¹⁶ Marine Scotland (2017) 'Marine Scotland Licensing Operations Team: Scoping Opinion for Seagreen Phase 1 Offshore Project'. Available at:
http://marine.gov.scot/sites/default/files/00524860_1.pdf (Accessed 01/09/2024)

¹⁷ Buckingham, L., Bogdanova, M.I., Green, J.A., Dunn, R.E., Wanless, S., Bennett, S., Bevan, R.M., Call, A., Canham, M., Corse, C.J., Harris, M.P., Heward, C.J., Jardine, D.C., Lennon, J., Parnaby, D., Redfern, C.P.F., Scott, L., Swann, R.L., Ward, R.M., Weston, E.D., Furness, R.W. and Daunt, F. (2022) 'Interspecific variation in non-breeding aggregation: a multi-colony tracking study of two sympatric seabirds'. Marine Ecology Progress Series 684: 181-197

Caledonia Offshore Wind Farm
5th Floor, Atria One
144 Morrison Street
Edinburgh
EH3 8EX

www.caledoniaoffshorewind.com

